

Research Article

Open Access



Pollen viability in natural populations of *Thymus capitatus* the south Al-Jabal Al-Akhdar area, Cyrenaica, Libya.

Ali, Ezzudin S., Khansa² A. Omar., Hesaen M. Mustafa³

¹Faculty of Agriculture, Omar Al-Mukhtar University, Libya.

² Faculty of Natural Resources, Omar Al-Mukhtar University, Libya.

³ Faculty of Natural Resources, Omar Al-Mukhtar University, Libya.

*Corresponding author: E-mail: Hussein.mohammad@omu.edu.ly
Environmental Sciences, Omar Al-mkhtar University elbayda, Libya

Received:
08. 05. 2025
Accepted:
31.08 .2025

Publish online: 31.08 .2025

الباحث الاول*: عز الدين، البستة، جامعة عمر المختار، ليبيا .
الباحث الثاني: خنساء، علوم البيئة، جامعة عمر المختار، ليبيا .
الباحث الثاني: مصطفى، حسين محمد علوم البيئة، جامعة عمر المختار، ليبيا .

Abstract: Viable pollen plays a crucial role in the process of seed formation and acts as the principal mechanism for gene transfer in outcrossing species such as thyme (*Thymus capitatus*). There is a lack of information on the pollen viability of natural populations of Libyan thyme that grow in the south El-Jabal El-Akhdar area. This research estimated pollen viability using acetocarmine staining method. It was possible to distinguish between viable pollen (non-aborted) and non-viable (aborted) pollen using acetocarmine stain. The viability of pollen percentage in *T. capitatus* accessions was found to be significantly different. The viable pollen recorded as highest was obtained from the white-flowered population (67.61%) followed by dotted white-flowered population (57.24%), while the lowest viable pollen was obtained from mosaic-flowered population (29.86%). The remaining two populations, violet-flowered and purple-flowered showed 55.73% and 42.38% respectively as pollen viability percentage. Variation was also observed in the pollen viability among different collection periods. These results suggested that pollen viability may be under both environmental and genetic control.

Keywords: Pollen viability; acetocarmine staining method; *Thymus capitatus*; El-Jabal El-Akhdar; Libya.

حيوية حبوب اللقاح في المجتمعات الطبيعية لنبات الزعتر البري *Thymus capitatus* في منطقة جنوب الجبل الأخضر، برقة، ليبيا.

المستخلص: حبوب اللقاح الخصبة والحية ضرورية لتكوين البذور، وتعمل بمثابة وسيلة أساسية لتدفق الجينات في الأنواع النباتية مفتوحة التلقيح مثل نبات الزعتر (*Thymus capitatus* L.). هناك نقص في المعلومات عن خصوبة حبوب اللقاح (حيوية حبوب اللقاح) في العشائر الطبيعية للزعتر البري، والذي ينتشر بصورة طبيعية في منطقة جنوب الجبل الأخضر. في هذا البحث قدرت حيوية حبوب اللقاح باستخدام صبغة الأسيتوكارمين. وقد بينت الدراسة فاعلية هذه الطريقة، فقد كان من الممكن التمييز بين حبوب اللقاح الخصبة (الحية) وحبوب اللقاح غير الخصبة (المجهضة). اشارت النتائج إلى أن الاختلافات في نسب حيوية حبوب اللقاح في العشائر الطبيعية للزعتر الليبي (*Thymus capitatus* L.) كانت ذات فروق معنوية. فقد سجلت عشيرة النباتات ذات الازهار البيضاء أعلى نسبة لحيوية حبوب اللقاح الخصبة (67.61%)، تليها مباشرة عشيرة النباتات ذات الازهار البيضاء المنقطعة (57.24%)، بالمقابل سجلت عشيرة النباتات ذات الازهار المبرقشة أقل نسبة لحيوية حبوب اللقاح الخصبة (29.86%). العشيرتين: عشيرة النباتات بنفسجية الازهار وعشيرة النباتات ارجوانية الازهار سجلت النسب 55.73% و 42.36% على التوالي. أيضاً بينت النتائج أن هناك تباين في حيوية حبوب اللقاح بين مختلف فترات التجميع. هذه النتائج تشير بأن حيوية حبوب اللقاح في عشائر الزعتر تخضع لفعل العوامل البيئية والعوامل الوراثية.

الكلمات المفتاحية: حيوية حبوب اللقاح، طريقة اختبار صبغة أسيتوكارمين، الزعتر البري، الجبل الأخضر - ليبيا.



INTRODUCTION

Effective pollination is essential for the successful development of fruits and seeds in the majority of plant species. Understanding pollen biology, particularly aspects related to pollen quality, is crucial for implementing strategies aimed at enhancing agricultural productivity (Shivanna, 2003). Pollen quality plays a significant role in the overall fitness of plants and marks a vital phase in their life cycle. It is commonly associated with pollen viability, which refers to the percentage of pollen grains that remain viable (Stanley and Linskens, 1974; Heslop-Harrison *et al.*, 1984).

The extensive body of literature concerning pollination ecology and pollen biology reveals a considerable diversity in the terminology and definitions associated with the functional capabilities of pollen. Viability is commonly defined as the "capacity to live, grow, germinate, or develop" (Lincoln *et al.*, 1982). Additionally, the concept of viability has been employed to refer to pollen grains that can germinate on the stigma (Morse, 1987; Niesenbaum, 1992), those that can germinate in vitro (Beardsell *et al.*, 1993; Lindgren *et al.*, 1995), as well as those that can absorb specific stains (Becker and Ewart, 1990; Nyman, 1992), and to indicate successful seed set following the process of pollination (Smith-Huerta and Vasek, 1984).

Research into the variability of pollen grains plays a crucial role in elucidating the mating systems of plants. The diversity of these mating systems significantly affects the variability and differentiation processes observed within populations of both wild and cultivated species (Grant, 1981; Richards, 1997). Furthermore, pollen viability is a key factor in addressing issues related to sterility, hybridization initiatives (Stone *et al.*, 1995), and the field of evolutionary ecology (Thomson *et al.*, 1994). This viability is particularly vital for understanding genetic variability in plants where cross-fertilization is more prevalent than self-fertilization (Divakara *et al.*, 2010). Generally, a direct correlation exists between pollen viability and the germination potential of pollen across numerous plant species (Stanley and Linskens, 1974). Numerous methodologies have been established to assess pollen viability, utilizing either the dimensions of pollen grains (Kelly *et al.*, 2002) or the dielectric characteristics of cellular membranes (Heidmann *et al.*, 2016; Heidmann and Di-Berardino, 2017). Nevertheless, even with the advent of innovative techniques, a significant portion of the scientific community continues to depend on traditional methods that necessitate only basic staining processes and a microscope for manual enumeration (Shivanna and Tandon, 2014).

Staining methods serve as effective tools for assessing the physiological condition of mature pollen grains. The appropriateness of a pollen viability test is contingent upon the specific species, as variations in optimal staining techniques have been documented (Rodriguez-Riano and Dafni, 2000). A range of staining methods has been utilized to evaluate the viability of pollen grains. Among these, acetocarmine, Alexander's stain, and aniline blue are the most commonly employed techniques for estimating pollen viability (Alexander, 1969; Migdałek *et al.*, 2014; Radovid *et al.*, 2017). The underlying principle of these staining reactions is based on the acidic nature of certain plant cell components, which exhibit a preference for basic dyes. The acetocarmine technique is extensively employed to assess pollen viability in both cultivated and wild plant species (Amkul *et al.*, 2016; Rathod *et al.*, 2018). Acetocarmine functions as a basic dye, with iron serving as a mordant. The nucleic acids, which possess a strong negative charge due to the phosphate groups present in the DNA structure, readily interact with basic dyes. Consequently, this method is utilized for swift evaluations of nuclear conditions and for in-depth investigations of chromosomes (Ali, 2008).

In Libya, Thyme (*Thymus capitatus*) is native to Al-Jabal Al-Akhdar area where it has been used for flavoring and medicinal purpose. Thyme is a candidate for large scale as a source of nectar for honey bees, and as an aromatic plant containing essential oils, which are important to the flavoring and medicinal applications (Ali and Mustafa, 2019).

Furthermore, an understanding of reproductive biology, particularly regarding pollen viability, is crucial for the conservation, management, and restoration of endangered species (Kuniyal *et al.*, 2003; Murugan *et al.*, 2006). Given that thyme (*Thymus capitatus*) holds significant medicinal and ecological value, it has garnered substantial research interest (Ali and Mustafa, 2019; Ali and Mustafa, 2021; Ali *et al.*, 2022). However, studies on the floral biology of *T. capitatus* have not been performed. Therefore, the objective of this research is to estimate the pollen viability of *T. capitatus* accessions (populations) that are growing in the south El-Jabal El-Akhdar area, Libya.

MATERIALS AND METHODS

Pollen samples of *Thymus capitatus* were collected from southern parts of El-Jabal El-Akhdar area, aiming to cover as much of the geographic range as possible. *Thymus capitatus* that growing in southern parts of Al-Jabal Al-Akhdar shows flower color polymorphism. There are five different accessions (phenotypes); white-flowered, dotted white-flowered, purple-flowered, violet-flowered and flowered, and mosaic-flowered accessions (Ali and Mustafa, 2019; Ali and Mustafa, 2021).

Thymus capitatus began to flower on in mid-july and generated pollen for approximately four weeks. Branches containing several inflorescences were harvested from 25 plants per accession (genotype) at three distinct periods during the pollination period: 1 week, 2 weeks, and 3 weeks. They were collected between 8:30 and 10:30 a.m. on sunny day mornings. The branches with inflorescences were placed in jars filled with tap water and maintained at room temperature (25 °C ±5 °C).

Pollen viability was determined by staining with acetocarmine (Ali and Bataw, 2014). Acetocarmine stains the pollen grain nuclei, and weakly stains the cytoplasm, and gives a good contrast between the grain and surrounding medium. The pollen nucleus is rich in chromatin material and live pollen stains pink to deep red with acetocarmine, whereas sterile (nonviable) pollen does not take any stain and thus remains almost white and transparent (McKellar and Quesenberry, 1992; Marutani, *et al.*, 1993). To prepare the acetocarmine stain, weigh 1 gram of carmine powder and dissolving it in 95 milliliters of glacial acetic acid. Subsequently, add distilled water to achieve a final volume of 100 milliliters. The solution should then be boiled, allowed to cool, filtered, and stored in a refrigerator. For the analysis of pollen viability, two to three drops of stain were applied to a slide, onto which pollen grains were subsequently sprinkled. A cover slip was then placed over the preparation, and the viability of the pollen was assessed after a duration of 5 to 10 minutes. For each of the five accessions (genotypes), at least 50 pollen grains were randomly counted in each visual field on the slide using a light microscope at a magnification of 100X. Each accession is analyzed using ten slides, from which images are obtained. That are red are identified as viable pollen. The average quantity of pollen for each accession is calculated, and the viability of the pollen is expressed as a percentage. The percentage of pollen viability is determined using the following formula:

$$\text{Pollen Viability (\%)} = \frac{\text{Number of stained pollen grains}}{\text{Total number of pollen grains on slide}} \times 100$$

Utilizing a completely randomized design (CRD), the experiment was set up. Analysis of variance (ANOVA) was performed on the experimental data. The significant F test (P 0.05) was used to determine the least significant differences.

RESULTS

According to the color of flower, there are five accessions that growing in south of Al-Jabal Al-Akhdar region, Cyrenaica, Libya: white-flowered, dotted white-flowered, violet-flowered, purple-flowered and mosaic-flowered accession (Ali and Mustafa, 2019; Ali and Mustafa, 2021; Ali, *et. al.* 2022).

In the current investigation, pollen viability was tested using acetocarmine stain which yielded a sufficiently clear result as shown in Figure 1. Since nonviable pollen didn't absorb the stain, which contrasted sharply with the red color of viable pollen, aceto-carmine stain was a dependable way to determine whether thyme pollen was viable (non-aborted) or nonviable (aborted); more significantly, the stain never colored aborted pollen (Fig. 1).




Figure: (1). Pollen grains of Libyan thyme (*Thymus capitatus*) tested by aceto-carmine stain;(a viable pollen, and non-viable pollen b).

Table 1 shows the results of pollen viability of the 5 accessions (genotypes) of Libyan thyme (*Thymus capitatus*) which growing in southern parts of Al-Jabal Al-Akhdar area, evaluated with aceto-carmine stain. The specimens examined generally produced well-formed pollen which had a high stain ability. The viability of pollen percentage in *T. capitatus* accessions was found to be significantly different (LSD=1.19 at $P \leq 0.05$) in the acetocarmine method. The viable pollen recorded as highest was obtained from the white-flowered population (67.61%) followed by dotted white-flowered population (57.24%), while the lowest viable pollen was obtained from the anthers of mosaic-flowered population (29.86%) in the acetocarmine stain test. The remaining two populations; violet-flowered and purple-flowered showed 55.73% and 42.38% respectively as pollen viability percentage (Table 1 and Fig. 2).

Table:(1). Viability percentages of pollen in libyan thyme (*Thymus capitatus*) accessions

Period of time	Means LSD = 2.08					Means LSD= 0.93
	White-flowered	Dotted white-flowered	Violet-flowered	Purple-flowered	Mosaic-flowered	
1week	78.08 ^a	70.83 ^b	64.45 ^c	52.15 ^f	38.30 ^h	60.74 ^A
2weeks	70.34 ^b	59.34 ^d	61.31 ^d	43.19 ^g	29.91 ^j	52.86 ^B
3weeks	54.48 ^e	41.60 ^g	43.48 ^g	34.89 ⁱ	21.18 ^k	38.10 ^C
Means LSD=1.19	67.61 ^A	57.24 ^B	55.73 ^C	42.38 ^D	29.86 ^E	

(a-b-c-d-e-f-g-h-i-j-k) These symbols indicate significant differences. If they differ, there is a significant difference. If the symbols are similar, there are no significant differences.

Variation was also observed in the pollen viability among different collection periods. The current results indicated that, pollen viability percentage of five accessions (populations) was significantly affected by the collection period (Table 1). The pollen viability percentage began with 60.74% for the first week, then recorded 52.86% for the second week. After three weeks, the pollen viability percentage sharply decreased to 38.10%. Those results were significantly different at the value of $L.S.D = 0.93$, $P \leq 0.05$.

Furthermore, the three collection periods within each single population (accession) showed different pollen viability (Fig. 2). For each accession, pollen viability was highest in the first collection period (the 1st week) and lowest in the third collection period (the 3rd week). The highest values of pollen viability occurred in the first period of collection (the 1st week) recording 78.08% (white-flowered), 70.83% (dotted white-flowered), 64.45% (violet-flowered), 52.15% (purple-flowered), and 38.30% (mosaic-flowered). In the second period of collection (the 2nd week), the values of pollen viability were 70.34% (white-flowered), 59.34% (dotted white-flowered), 61.31% (violet-flowered), 43.19% (purple-flowered) and 29.91% (mosaic-flowered). The lowest percentages of pollen viability occurred after three weeks (the 3rd period of collection) recording 54.48% (white-flowered), 41.61% (dotted white-flowered), 43.48% (violet-flowered), 34.89.19% (purple-flowered) and 21.18% (mosaic-flowered). Those results were significantly different at the value of $L.S.D = 0.93$, $P \leq 0.05$.




Figure: (2). Viability percentages of pollen during the three collection periods in Libyan thyme accessions

DISCUSSION

The results clearly showed that pollen viability using acetocarmine stain was effective in identifying fertile pollen grains, thus obtaining an early indication of pollen quality. According to several previous studies (Thomson, 1989; Dafni and Firmage, 2000; Bots and Mariani, 2005; Ali and Batao, 2014), pollen viability is a crucial indicator of pollen quality. The viability of pollen grains, which can be evaluated using a variety of techniques, including staining with non-vital dyes, is typ-

ically a prerequisite for successful seed set (Dafni and Firmage, 2000; Dafni *et al.*, 2010; Guoren He *et al.*, 2017). One typical method for assessing pollen viability is acetocarmine staining (Migdalek *et al.*, 2014; Maryam *et al.*, 2015; Vijayakumar *et al.*, 2018).

The findings of the current research indicated that the acetocarmine staining technique effectively distinguishes between viable (non-aborted) and non-viable (aborted) pollen grains in populations of Libyan thyme (*T. capitatus*). Also, the results corresponded well with that of Tiwari *et al.* (2014) who evaluated pollen viability in members of *Euphorbiaceae* family using several dyes and concluded that acetocarmine staining method was efficient for distinction of viable and nonviable pollen grains and provided clear results. Thus, acetocarmine staining method can be recommended for testing pollen viability. Results were also in agreement with other studies indicating that the use of different colorants to test pollen viability may give comparable results (Parfitt and Ganeshan, 1989; Ali and Bataw, 2014).

Temperature, moisture, genotypic variations, plant vigor, physiological stage, and flower age all have a significant impact on pollen survival (Adhikari and Campbell, 1998; Ali and Bataw, 2014; Shivashankara *et al.*, 2019; Parashuram *et al.*, 2021). In the current investigation, the rates of pollen viability were significantly differed among all accessions (genotypes) of libyan thyme. The highest viable pollen rate was recorded from the white-flowered population (67.61%), while the lowest was obtained from the mosaic-flowered population (29.86%). Analysis of variance confirmed the significance of these differences, suggesting high variation in pollen viability among accessions with a 95% confidence interval. The different percentage of pollen viability can be attributed to the genotype, since male inflorescences were sampled from the same site and at the same physiological stage. Genotypic differences among accessions in others plant species for pollen viability have been previously reported (Sharafi, 2011; Ali and Bataw, 2014; Lankinen *et al.*, 2018).

Pollen viability is a crucial factor that can restrict seed production (Fritz and Lukaszewski, 1989; Dafni and Firmage, 2000). The findings of the current study revealed that in natural populations of Libyan thyme, pollen viability can significantly decrease after a period of three weeks. Consequently, if pollen arrives at the pistil more than three weeks later, successful pollination may not occur. In general, percentages of pollen viability were decreased with time in all accessions of Libyan thyme. Similar results were reported in many previous studies indicating that pollen viability decreased with time (Beyhan and Serdar, 2008; Ali and Bataw, 2014; Slomka, *et al.*, 2014; Mantiquilla *et al.*, 2018; Robles-Gonzalez *et al.* 2019).

CONCLUSION

The main conclusions of the experimental work should be presented. The contribution of the work to the scientific community and its economic implications should be emphasized.

REFERENCES

- Adhikari K. and Campbell C. (1998). *In vitro* germination and viability of buckwheat (*Fagopyrum esculentum* Moench) pollen. Euphytica 102(1): 87–92. DOI: [10.1023/A:1018393425407](https://doi.org/10.1023/A:1018393425407)
- Alexander MP (1969). Differential staining of aborted and non-aborted pollen. Stain Technology 44: 117-122. DOI: [10.3109/10520296909063335](https://doi.org/10.3109/10520296909063335)

- Ali, E. S. (2008). Behavior of chromosomes during meiosis in wild olive. Journal of Agricultural Research, Kafrelsheikh University, 34(3): 808-819. DOI: <https://doi.org/10.37375/bsj.vi17.2751>
- Ali, E. S. and Bataw, M. (2014). Pollen grains viability in *Thapsia garganica* from Al-jabal Al-akhder region, Libya. Al-Mukhtar Journal of Sciences, 29(1):140-148. (in Arabic, with English abstract).
- Ali, E. S. and Mustafa, H. M. (2019). Polymorphism in thyme (*Thymus capitatus*) at southern region of El-Jabal El-Akhdar, Libya. Bayan Scientific journal, 4:47-62. (in Arabic, with English abstract).
- Ali, E. S. and Mustafa, H. M. (2021). Isoenzyme diversity for peroxidase among Libyan thyme (*Thymus capitatus*) populations. Bayan Scientific journal, 9:658-664.
- Ali, E. S., Mustafa H. M. and K. A. Omaar (2022). Phylogenetic Analysis of Libyan Thyme (*Thymus Capitatus*) Inferred from The Morphological Traits. Al-Mukhtar Journal of Sciences 37 (4): 385-393. DOI: [10.54172/mjsc.v37i4.978](https://doi.org/10.54172/mjsc.v37i4.978)
- Amkul, K., Panngam, M., Tanya, P., Srinives, P. and Laosatit, K. (2016). Pollen viability and seed set of interspecific hybrids between *Jatropha curcas* × *Jatropha integerrima*. Genomics and Genetics, 9(1): 50-55. DOI: [10.14456/gag.2016.7](https://doi.org/10.14456/gag.2016.7)
- Beardsell, D. V., Knox, R. B. and Williams, E. G. (1993). Breeding system and reproductive success of *Thryptomene calycina* (Myrtaceae). Australian Journal of Botany, 41:333-353.
- Becker, W. A. and Ewart, L. C. (1990). Pollination, seed set and pollen tube growth investigation in *Viola redata* L. Acta Horticulture, 272:33-36. DOI: [10.17660/ActaHortic.1990.272.3](https://doi.org/10.17660/ActaHortic.1990.272.3)
- Beyhan N. and Serdar U. (2008). Assessment of pollen viability and germinability in some European chestnut genotypes (*Castanea sativa* L.). Hort. Sci. 35(4): 171–178. DOI: [10.17221/23/2008-HORTSCI](https://doi.org/10.17221/23/2008-HORTSCI)
- Bots M. and Mariani C. (2005). Pollen viability in the field. Radboud Universiteit Nijmegen. Report COGEM. p. 13–15.
- Dafni A. and Firmage D. (2000). Pollen viability and longevity: Practical, ecological and evolutionary implications. Plant Syst. Evol. 222(1–4): 113–132. <https://doi.org/10.1007/BF00984098>
- Dafni, A., Kevan P., Gross L. and K. Goka (2010). *Bombus terrestris*, pollinator, invasive and pest: An assessment of problems associated with its widespread introductions for commercial purposes. Appl. Entomol. Zool. 45 (1): 101–113. DOI: [10.1303/aez.2010.101](https://doi.org/10.1303/aez.2010.101)
- Divakara, B. N., Upadhyaya, H. D., Wani, S. P., Laxmipathi-Gowda, C. L. (2010). Biology and genetic improvement of *Jatropha curcas* L.: a review. Appl. Energy, 87: 732-742. <https://doi.org/10.1016/j.apenergy.2009.07.013>
- Firtz S.E. and Lukaszewski A.J. (1989). Pollen longevity in wheat, rye and triticale, Plant Breeding, 102:31-34. <https://doi.org/10.1111/j.1439-0523.1989.tb00311.x>

Grant, V. (1981). Plant Speciation. 2nd edition. New York, Columbia Univ. Press.

Guoren He, Fengrong Hu, Jun Ming, Chun Liu and Suxia Yua (2017). Pollen viability and stigma receptivity in *Lilium* during anthesis *Euphytica*, (1): 213-231. DOI: [10.1007/s10681-017-2019-9](https://doi.org/10.1007/s10681-017-2019-9)

Heidmann, I., and Di-Berardino, M. (2017). Impedance flow cytometry as a tool to analyze microspore and pollen quality. *Methods Mol. Biol.*, 1669: 339–354. DOI: [10.1007/978-1-4939-7286-9_25](https://doi.org/10.1007/978-1-4939-7286-9_25)

- Heidmann, I., Schade-Kampmann, G., Lambalk, J., Ottiger, M., and Di Berardino, M. (2016). Impedance flow cytometry: a novel technique in pollen analysis. *PLoS One*. <https://doi.org/10.1371/journal.pone.0165531>

Heslop-Harrison, J., Heslop-Harrison, Y. and Shivanna, K. R. (1984). The evaluation of pollen quality, and a further appraisal of the fluorochromatic (FCR) test procedure. *Theor. Appl. Genet.*, 67: 367-375. doi.org/10.1007/BF00272876

- Kelly, J. K., Rasch A. and Rasch, S. Rasch. (2002). A method to estimate pollen viability from pollen size variation. *American Journal of Botany*, 89: 1021–1023. DOI: [10.3732/ajb.89.6.1021](https://doi.org/10.3732/ajb.89.6.1021)

Kuniyal, C. P., Rajsekaran, C. Prasad, P. and Bhadula, S. K. (2003). Multiplication of *Aconitum atrox* (Bruhl) Muk., via tuber segments. *Int. Plant Genet. Resour. Newsletter*, 135:59-62.

Lankinen A., Sandra A. M., Lindstrom and Tina D'Hertefeldt (2018). Variable pollen viability and effects of pollen load size on components of seed set in cultivars and feral populations of oilseed rape. <https://doi.org/10.1371/journal.pone.0204407>.

Lincoln, R. J., Boxshall, G. A. and Clark, P. F. (1982). A dictionary of ecology, evolution and systematics. Cambridge University Press.

Lingren, D., Paule, L., Xihuan, S., Yazdani, R., Segerstrom, U., Wallen, J. and Lijdebro, M. L. (1995). Can viable pollen carry Scots pine genes over long distances? *Grana*, 34:64-69. DOI: [10.17660/ActaHortic.2001.561.13](https://doi.org/10.17660/ActaHortic.2001.561.13)

Mantiquilla J. A., Elumba M., Adtoon1 J. A., Abad R. G., Cyrose S., Millado S., and Gilda C. R. (2018). *In vitro* germination and viability testing of Nipa(*Nypa fruticans* Wurmb.) pollen under different storage conditions. *Philippine Journal of Science* 147 (4): 617-627.

Maryam, M, Jafar, J., Bilques, F., Salman, H., Summar, A., Naaqvi, N., Rashid, A., and Iqar, A. Khan (2015). Evaluation of pollen viability in date palm cultivars under different storage temperatures. *Pak. J. Bot.*, 47(1): 377-381.

Marutani, M., Sheffer, R. D. and Kameto, H. (1993). Cytological analysis of *Arthuriium andraenum* (Araceae), its related taxa and their hybrids. *American Journal of Botany*, 80:93-103. <https://doi.org/10.1002/J.1537-2197.1993.TB13772.X>

McKellar, M. A and Quesenberry, K. H. (1992). Chromosome pairing and pollen viability in *Desmodium ovalifolium* Wall x *Desmodium heterocarpon* (L.) DC hybrids. *Australian Journal of Botany*, 40:243-247. <https://doi.org/10.1071/BT9920243>

- Migdałek, G., Kolczyk, J., Pliszko, A., Kościńska-Pająk, M. and Słomka, A. (2014). Reduced pollen viability and achene development in *Solidago* × *niederederi* Khek from Poland. *Acta Soc. Bot. Pol.*, 83(3):251–255. [DOI: 10.5586/asbp.2014.025](https://doi.org/10.5586/asbp.2014.025)
- Morse, D. H. (1987). Roles of pollen and ovary age in follicle production of the common milkweed *Asclepias syriaca*. *American Journal of Botany*, 72: 527-536.
- Murugan, R., Shivanna, K. R. and Rao, R. R. (2006). Pollination biology of *Aristolochia tagala*, a rare species of medicinal importance. *Curr. Sci.*, 91(6):795-798.
- Niesenbaum, R. A. (1992). Sex ratio, components of reproduction and pollen deposition in *Linderabenzoin* (Lauraceae). *American Journal of Botany*, 79(5):495-500. <https://doi.org/10.1002/j.1537-2197.1992.tb14584.x>
- Nyman, Y. (1992). Pollination mechanisms in six *Campanula* species (Campanulaceae). *Pl. Syst. Evol.*, 181:97-108. <https://doi.org/10.1007/BF00937589>
- Parfitt D.E. and Ganeshan S. (1989). Comparison of procedures for estimating viability of *Prunus* pollen. *HortScience* 24:354–356
- Parashuram, M., Pavan Kumar, P., Raghavendra, G., Shirol, A., Bapuraygouda, P, Jameel M. and Shankar, M. (2021). Assessment of pollen viability and *in vitro* pollen germination in *Nerium* cultivars (*Nerium oleander* L.). *International Journal of Chemical Studies*, 9(2): 724-728. [DOI: https://doi.org/10.22271/chemi.2021.v9.i2k.11898](https://doi.org/10.22271/chemi.2021.v9.i2k.11898)
- Radovid, A., Nikolid, D., Milatovid, D., Rakonjac, V. and Bakid, I. (2017). Pollen viability in some Quince cultivars. 3rd International Symposium for Agriculture and Food (ISAF). 72(1):68-71. [DOI: 10.55302/JAFES18721068r](https://doi.org/10.55302/JAFES18721068r)
- Rathod V., Behera T., Munshi A., Durgesh K., Krishnan B. and Sharma N. (2018). Pollen viability and *in vitro* pollen germination studies in *Momordica* species and their intra and inter-specific hybrids. *International Journal of Chemical Studies*, 6(6): 32-40.
- Richards, A. J. (1997). *Plant Breeding Systems*. Chapman and Hall, London.
- Robles-González, M., Cortez-Arroyo, A., Carrillo-Medrano, S. and Medina-Urrutia, V. (2019). Viability and longevity of pollen in Mexican lemon genotypes estimated by *in vitro* germination. *Revista Mexicana Ciencias Agrícolas*, 10 (4):911-920. <https://doi.org/10.29312/re-mexca.v10i4.1780>
- Rodriguez-Riano, T. and Dafni, A. (2000). A new procedure to assess pollen viability. *Sex. Pl. Repr.*, 12:241-244.
- Sharafi, Y. (2011). Investigation on pollen viability and longevity in *Malus pumila* L., *Pyrus communis* L., and *Cydonia oblonga* L., *in vitro*. *Journal of Medicinal Plants Research*, 5(11): 2232-2236.
- Shivanna, K. R. (2003). *Pollen Biology and Biotechnology*. Science Publishers, Inc., USA, pp. 181–182.
- Shivanna, K. R. and Tandon, R. (2014). *Reproductive Ecology of Flowering Plants: A Manual*; Springer. New Delhi, India. <https://doi.org/10.1007/978-81-322-2003-9>

- Shivashankara, K. S., Geetha, G. A. and Roy, T. K. (2019). Metabolite profiling in Mango (*Mangifera indica* L.) pollen grains in relation to viability. J. Hortl. Sci., 14(1) : 33-42.
- Słomka A, Bohdanowicz J, Poznańska E, Kwiatkowska M, Pilarska M and Struebig M (2014). Usefulness and limitations of pollen characters in environmental studies based on *Viola* L. species (sect. *Melanium* Ging.). Mod Phytomorphology, 5:31–34.
- Smith-Huerta, N. L. and Vasek, F. C. (1984). Pollen longevity and stigma pre-emption in *Clarkia*. American Journal of Botany, 71(9):1183-1191.
- Stanley, R. and Linskens, H. (1974). Pollen. Biology, biochemistry, management. -Springer. Berlin, Heidelberg.
- Stone, J. L., Thomson, J. D. and Dent, A. S. J. (1995). Assessment of pollen viability in hand-pollination experiments: A review. American Journal of Botany, 82:1186–1197. <https://doi.org/10.1002/j.1537-2197.1995.tb11591.x>
- Thomson, JD (1989). Germination schedules of pollen grains: implications for pollen selection. Evolution; 43: 220-223. <https://doi.org/10.1111/j. PMID: 28568494>
- Thomson, JD., Rigney, L. P, Karoly, K. M. and Thomson, B. A. (1994). Pollen viability, vigor, and competitive ability in *Erythronium grandiflorum* (Liliaceae). American Journal of Botany, 81:1257–1266.
- Tiwari A., Kamble RB., and Chaturvedi A. (2014). Pollen biology of some members of Euphorbiaceae family. Asian J Plant Sci Res, 4: 8–14.
- Vijayakumar, R., TK B., Munshi, A. Kumar, D., G. Jat, Boopala Krishnan, G. and N. Sharma (2018). Pollen viability and *in vitro* pollen germination studies in *Momordica* species and their intra and interspecific hybrids. International Journal of Chemical Studies, 6(6): 32-40.