



Phytochemical Screening and Antibacterial Activity of *Moringaoleifera* Leaf Extract Western Libya

Safa E. Rahuma¹, Munira M. Ashrif^{*2}

***Corresponding author:** m.ashrif@zu.edu.ly, Department of Chemistry, Faculty of Science, University of Zawia, Zawia, Libya.

Second Author: safaeltabeb@gmail.com Department of Chemistry, Faculty of Education Zaltan, Sabratha University, Zaltan, Libya.

Received:
20 December 2022

Accepted:
24 October 2023

Publish online:
31 December 2023

Abstract

Moringa oleifera (*M. oleifera*) is one of the Moringaceae family. *M. oleifera* plant parts retain a precious pharmacological effect. Its leaves are utilized commonly because of having enormous nutrients as nutraceuticals. Moreover, its compresses various effects such as biological and antimicrobial effect as antibacterial or antifungal. Additionally, *M. oleifera* investigationin was carried out either in vivo or in vitro to conclude that their leaves component has pharmacological effect. This current study had confirmed therapeutic effect of *Moringa Oleifera* leaf extract and its antibacterial activity. There is critical need for more advanced research about use and pharmaceutical effects of *Moringa Oleifera* leaves in improvement of various drugs and useful foods. This study aimed to investigate phytochemical ingredients and antibacterial effect of *Moringa Oleifera* leaf extract and the uses of *Moringa Oleifera* leaf extract either traditionally or pharmacologically concluded.

Keywords: *Moringaoleifera*, Antibacterial activity, Phytochemical screening, Libya.

INTRODUCTION

The drumstick tree, or *Moringaoleifera* Lam., is a member of the Moringaceae family and is extensively dispersed in various regions, including tropical and desert nations, as well as in India and Africa. It was primarily used for food and medicine (Al_husnan & Alkahtani, 2016). *Moringaoleifera* has numerous qualities that may be used for food, medicine, and other industrial and home applications (Falowo et al., 2018; Padayachee & Baijnath, 2020), particularly its leaves, which may be eaten raw, roasted, or dried and kept for a long time without losing any of their nutritional value. In addition to using the leaves for food and feed, they may also be used as functional foods or nutraceuticals since they naturally contain phytochemicals such as flavonoids, carotenoids, and glucosinolates (Amaglo et al., 2010; Sultana et al., 2009). The main phytochemicals found in moringa leaves are astragalin, isoquercetin, and crypto-chlorogenic acid, which have anti-oxidant, anti-hypertension, and anti-inflammation characteristics (Verma et al., 2009; Vongsak et al., 2012).

Based on the bioactive components and their antioxidant activity, several in vitro experiments have largely supported the biological activity of these plant extracts (Leone et al., 2015; Sultana et al., 2009). Its strong antioxidant properties are principally brought on by its high phenolic content. Due to these therapeutic advantages, several pharmaceutical preparations from this plant have been manufactured and sold in both the Indian and global markets.(Mehra et al., 2020; Sahakitpichan et al., 2011) Different sections of *M. oleifera* have antibacterial potentials that can be used to purify



The Author(s) 2023 This article is distributed under the terms of the *Creative Commons Attribution-NonCommercial 4.0 International License* (<http://creativecommons.org/licenses/by-nc/4.0/>) (<http://creativecommons.org/licenses/by-nc/4.0/>)), which permits unrestricted use, distribution, and reproduction in any medium, *for non-commercial purposes only*, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

water, eliminate biofilm threats, and eradicate pathogenic microbes. The long-term use of *M. oleifera* and its ethno-pharmacological characteristics have proven its safety. *M. oleifera* has also been employed for bio-enhancement and as nanoparticles in medication administration in addition to its antibacterial action (Arora et al., 2013). Fidrianny et al. study mentioned that *M. oleifera* is commonly named Sanjana and drumstick. The name Moringa originated from murungai; a Tamil word that means twisted pod (Dhakad et al., 2019; Fidrianny et al., 2021) study illustrated that the cultivation of Moringa has different reasons according to its nutritional content or medicinal uses (Djemoui et al., 2019). *M. oleifera* has medicinal effects documented by (Djemoui et al., 2019). study, such as hypoglycemic effect, antioxidant, antibacterial, antifungal, and wound healing due to anti-inflammatory activity (Djemoui et al., 2019). (Amaglo et al., 2010) reported that the leaves, in particular, can be used in a salad, roasted, or kept as dry powder for a prolonged time without losing nutritive constituents (Amaglo et al., 2010).

Moringa leaves were reported by (Maheshwari et al., 2014). to have elevated amounts of protein that could be utilized as food. Also, they mentioned that *Moringa oleifera* contains a high amount of provitamin A.(Tshabalala et al., 2020). traced the source of the cardiovascular benefit and body energizing to the omega-3 and omega-6 polyunsaturated fatty acid content of Moringa. It can be used for obese patients due to low calorific value (Kashyap et al., 2022; Maheshwari et al., 2014)

They also illustrated calcium contents and digestible protein that is higher than milk. (Kasolo et al., 2010). mentioned moringa leaves' high content of potassium, magnesium, and copper. Iron content is 28 times of spinach, proved (Fuglie, 1999; Kasolo et al., 2010). Interestingly, Mbikay approved that fat-soluble vitamins like D and E and water-soluble like folic acid, nicotinic acid, and vitamin care also exist in *M. oleifera* (Mbikay, 2012). Stevens et al. mentioned that leaves are not toxic when taken in large amounts, so it is found to be safer and healthier for consumption (Stevens et al., 2013). Bacterial species develop drug resistance to common antibacterial agents. So, medicinal plants such as Moringa leaves could be used as alternative treatments for infectious diseases (Luqman et al., 2012).

Rios & Recio's studies reported the need for more studies on the antibacterial activity of essential oils or compounds like alkaloids, flavonoids, diterpenes, and triterpenes to justify the use of medicinal plants (Rios & Recio, 2005), while Abdallah mentioned that metabolites of *Moringa oleifera* leaves, such as alkaloids, flavonoids, and saponins have wonderful pharmacological effects as antibacterial and anti-microbial properties and interestingly as anticancer on (Berkovich et al., 2013). study (Abdallah, 2011; Berkovich et al., 2013).

MATERIALS AND METHODS

Collection and Identification of Plant Material

Moringaoleifera tree leaves were collected. The leaves were washed under distilled water for dust and foreign particle removal. After that, the leaves were air-dried in an oven at 60°C. The dried leaves were added to a mixer to be ground into fine powder. The fine powder was stored in a refrigerator to avoid heat and light exposure and kept for further use.

Preparation of Plant Material

About 10g of the fine powdered *Moringaoleifera* leaves were suspended in 100 mL of ethanol and shaken well. The suspensions were kept at room temperature for 24 hours. The suspensions were then filtered. The solvent was removed by heat treatment at 80 °C to concentrate the extract. The extracts were stored in a refrigerator at 4°C for further analysis (Abadallah & Ali, 2019).

Phytochemical Screening

The collected plant extracts were subjected to qualitative phytochemical analysis for identification of various bioactive chemical constituents (alkaloids, volatile oils, saponins, flavonoids, steroids, tannins, and terpenoids) conducted using standard laboratory methods as described by (Edeoga et al., 2006; Sofowora, 1993).

Test Isolates

Clinical *Staphylococcus aureus*, *Streptococcus pyogenes*, and *Pseudomonas aeruginosa* isolates were collected for testing the antibacterial activity of *Moringaoleifera*. As described by Cheesbrough, gram staining and microbiological analysis were used in recognizing the isolates by utilizing differential media and biochemical tests.

Antibacterial activity testing

0.1 ml of each organism culture was inoculated onto the plates' surface by a sterile swab stick. The plates were incubated for 24 hours at 37°C, after which the inhibition zones were calculated using a meter ruler.

All the experiments were repeated three times, and the mean was recorded.

RESULTS

Phytochemical Screen

1. Tannins: A green color and gelatinous precipitate resulted upon adding 1 ml of 1% Lead acetate to 1 ml of the moringa leaf extract in a test tube. This result confirmed the existence of tannins.
2. Saponins: Durable white precipitate resulted when 1-3 ml of mercury chloride was added to 5 ml of leaf extract which was judged as an indicator of saponins existence.
3. Flavonoids: A yellow shade that signified flavonoids resulted when moringa leaf extract was added to diluted ammonia and concentrated sulphuric acid.
4. Terpenoids (Salkowskitest): No brown color resulted when 1 ml of leaf extract was added to chloroform, one drop of concentrated sulphuric acid, and one drop of anhydrous acetic acid.
5. Resins: Turbidity resulted when the leaf extract was added to the ethyl alcohol (95%) solution
6. Volatile oils: A shiny pink layer on filter paper formed after UV light exposure after the filtration of leaf extract, indicating volatile oils' content.
7. Alkaloids: A grey color confirmed alkaloids' presence when adding leaf extract to 2 ml of Marquis Reagents.
8. Glycosides: A few drops of Fehling reagent were added to 1 ml of leaf extract.

(Table 1) lists the phytochemical components of moringa leaf extracts.

Table: (1). Illustrate the phytochemical constituent of moringa leaf extract.

Phytochemical constitutes	Result
Alkaloids	++
Saponins	+
Tannins	+++
Flavonoids	+
Terpenoids	-
Resins	+++
Volatile oils	+++
Glycosides	++

Symbol:+=Existence of phytochemical, -=Absence of phytochemical. (+++)high, (++) medium, (+) poor, (-)no found

The results indicate the presence of saponins, resins, tannins, alkaloids, flavonoids, glycosides and volatile oils, and the absence of terpenoids.

Antibacterial Activity:

The antibacterial activity of *Moringa Oleifera* leaf extract is summarized in (Table 2).

The inhibition zones emphasized that it depends on the type of bacteria and concentration of *Moringa Oleifera* leaf extract. The highest zone of inhibition is demonstrated by *Pseudomonas aeruginosa* (17 mm) in concentrated extract and (14mm) in diluted extract.

Interestingly, the case of *Staphylococcus aureus* and Streptococcus pyogenes reported zero results which indicates their negative results and the antibacterial activity of *Moringa Oleifera* leaf extract.

Table: (2). Demonstrate the antibacterial activity of *MoringaOleifera* leaves extract

Name		Result
Staphylococcusaureus	Concentrated	0
	Diluted	0
Streptococcus pyogenes	Concentrated	0
	Diluted	0
Pseudomonasaeruginosa	Concentrated	17
	Diluted	14

DISCUSSION

Chhetri *et al.* study reported that bioactive constituents can be taken from plant parts like seeds, roots, stems, leaves, and flowers (Chhetri *et al.*, 2008). The results of our study showed the existence of bioactive components in moringa leaf extracts. Our study revealed the presence of alkaloids, volatile oils, saponins, resins, etc., in the ethanol extract of the leaves as in (Table1). The outcome of the current study is in a parallel line with previous studies that screened the phytochemical composition of moringa leaves. Moringa leaves were proven to include numerous phytochemicals (Edeoga *et al.*, 2006; Madziga *et al.*, 2010).

The phytochemicals have remarkable pharmaceutical effects that are used in drugs for treating sclerosis, diabetes, skin antiseptic, diarrhea, colitis, and cancer.(Aboaba *et al.*, 2011; Madziga *et al.*, 2010). studies illustrated that Alkaloids, for example, are extensively used as anticancer agents, aesthetics, and central nervous stimulants. Alkaloids also have metabolic effects and control development in living systems (Aboaba *et al.*, 2011; Madziga *et al.*, 2010).(Mir *et al.*, 2013). mentioned the role of saponins as one of the phytochemicals in cholesterol lowering. Tannins containing plants are commonly used as mouthwashes, and eye washes, as mentioned by (Abadallah & Ali, 2019; Mir *et al.*, 2013).

They also mentioned that terpenoids are effective as antimicrobial, anti-inflammatory, and immunomodulatory agents and in the case of cancer. (Abadallah & Ali, 2019) explained the result of moringa leaf extracts' antibacterial activity achieved in our study against the clinical isolates of *Staphylococcus aureus*, Streptococcus pyogenes, and pseudomonas (Abadallah & Ali, 2019).

The outcome of our study was in line with (Saadabi & Zaid, 2011) results that reported that the extracts of *moringaoleifera* have an impeding effect on pathogenic bacteria like *Pseudomonas aeruginosain* and *Escherichia coli* (Saadabi & Zaid, 2011). The result of this study concludes that ethanol extracts of *moringaoleifera* showed a maximum zone of inhibition (17 mm) against *Pseudomo-*

nas aeruginosainas in (Table 2). This conclusion also describes the study by (Al-Bakri & Afifi, 2007) who mentioned moringa leaf extract constraining the growth of *E. coli*, *Klebsiella pneumoniae*, and *P.aeruginosa* (Al-Bakri & Afifi, 2007). Numerous investigations have shown that *M. oleifera* leaves have antibacterial properties. As an illustration, a study on plant extracts prepared in different solvents found the ether extract of leaves of *M. oleifera* was the most active ingredient against the ambient and clinical isolates of *Proteus mirabilis*, a well-known cause of urinary tract infections (Arun & Rao., 2001). Several studies on the antifungal activity of *M. oleifera* leaves against *Trichophytonrubrum*, *T. mentagrophytes*, *Epidermophytonfloccosum*, and *Microsporum* documented that the crude extracts as well as essential oils from *M. oleifera* leaves reveal antifungal activity (Abalaka et al., 2012; Chuang et al., 2007). (Patel et al., 2014) study showed *M. oleifera* leaf extracts in ethanolic and aqueous forms were effective against the yeast *Saccharomyces cerevisiae* and the fungus *Candida tropicalis*, but not against *C. albicans*. Little is known about the antibacterial activity of *M. oleifera* roots. Nevertheless, a few investigations have revealed this plant's roots to have anti-microbial properties(Patel et al., 2014). Additionally, in vitro tests on several *M. oleifera* root bark extracts against *S. aureus*, *E. coli*, *Salmonella gallinarum*, *P. aeruginosa*, and others showed that ethyl acetate and acetone extracts have the highest activity in comparison to other solvents (Raj et al., 2011). Similar to this, only a small amount of research—a few early reports—has been done on the antibacterial properties of *M. oleifera's* stem bark. Studies demonstrate the existence of active organic extracts with variable degrees of activity in *M. oleifera* root extract's antibacterial efficacy against several human diseases (Chhetri et al., 2008).

CONCLUSION

The current study showed that *moringa oleifera* leaf extracts have an antibacterial activity that impedes bacterial growth. The results express that ethanol extract of *moringa oleifera* leaf is effective against bacterial strains.

The antibacterial effect may be due to bioactive constituents such as alkaloids, saponins, flavonoids, and tannins. Therefore, the study results emphasized the *moringa oleifera* leaf components and its therapeutic prospects as an antibacterial agent.

Duality of interest: In relation to this paper's publication, the authors state that they have no conflicts of interest.

Author contributions: Practical experimentation was conducted with Safa E. Rahuma assistance. Munira M. Ashrif contributed in writing, reviewing, and debating the paper.

Funding: No specific grant from a public or commercial funding agency was received for this research.

REFERENCES

- Abadallah, M., & Ali, M. (2019). Antibacterial activity of *Moringa oleifera* leaf extracts against bacteria isolated from patients attending general Sani Abacha specialist hospital damaturu. *J. Allied Pharm. Sci*, 1, 61-66.
- Abalaka, M., Daniyan, S., Oyeleke, S., & Adeyemo, S. (2012). The antibacterial evaluation of *Moringa oleifera* leaf extracts on selected bacterial pathogens. *Journal of Microbiology research*, 2(2), 1-4.

- Abdallah, E. M. (2011). Plants: An alternative source for antimicrobials. *Journal of applied pharmaceutical science*(Issue), 16-20.
- Aboaba, O. O., Ezeh, A. R., & Anabuiké, C. L. (2011). Antimicrobial activities of some Nigerian spices on some pathogens. *Agriculture and Biology Journal of North America*, 2(8), 1187-1193.
- Al-Bakri, A. G., & Afifi, F. U. (2007). Evaluation of antimicrobial activity of selected plant extracts by rapid XTT colorimetry and bacterial enumeration. *Journal of Microbiological Methods*, 68(1), 19-25.
- Al_husnan, L. A., & Alkahtani, M. D. (2016). Impact of Moringa aqueous extract on pathogenic bacteria and fungi in vitro. *Annals of Agricultural Sciences*, 61(2), 247-250.
- Amaglo, N. K., Bennett, R. N., Curto, R. B. L., Rosa, E. A., Turco, V. L., Giuffrida, A., Curto, A. L., Crea, F., & Timpo, G. M. (2010). Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* L., grown in Ghana. *Food Chemistry*, 122(4), 1047-1054.
- Arora, D. S., Onsare, J. G., & Kaur, H. (2013). Bioprospecting of *Moringa* (Moringaceae): microbiological perspective. *Journal of pharmacognosy and phytochemistry*, 1(6), 193-215.
- Arun T, Rao PCH. (2011). Phytochemical screening and antibacterial activity of *Moringa oleifera* Lam. against *Proteus mirabilis* from urinary tract infected patients. *International Journal of PharmTech Research*;3(4):2118–23.
- Berkovich, L., Earon, G., Ron, I., Rimmon, A., Vexler, A., & Lev-Ari, S. (2013). *Moringa Oleifera* aqueous leaf extract down-regulates nuclear factor-kappaB and increases cytotoxic effect of chemotherapy in pancreatic cancer cells. *BMC complementary and alternative medicine*, 13, 1-7.
- Chhetri, H. P., Yogol, N. S., Sherchan, J., Anupa, K., Mansoor, S., & Thapa, P. (2008). Phytochemical and antimicrobial evaluations of some medicinal plants of Nepal. *Kathmandu university journal of science, engineering and technology*, 4(1), 49-54.
- Chuang, P.-H., Lee, C.-W., Chou, J.-Y., Murugan, M., Shieh, B.-J., & Chen, H.-M. (2007). Anti-fungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresource technology*, 98(1), 232-236.
- Dhakad, A. K., Ikram, M., Sharma, S., Khan, S., Pandey, V. V., & Singh, A. (2019). Biological, nutritional, and therapeutic significance of *Moringa oleifera* Lam. *Phytotherapy Research*, 33(11), 2870-2903.
- Djemoui, D., Saidi, M., Rahmani, Z., & Djemoui, A. (2019). Influence of phenolic compounds on antioxidant capacity of leaves extracts of *Moringa oleifera* from Tamanrasset region. *Journal of Fundamental and Applied Sciences*, 11(1), 280-293.
- Edeoga, H., Omosun, G., & Uche, L. (2006). Chemical composition of *Hyptis suaveolens* and *Ocimum gratissimum* hybrids from Nigeria. *African journal of Biotechnology*, 5(10).

- Falowo, A. B., Mukumbo, F. E., Idamokoro, E. M., Lorenzo, J. M., Afolayan, A. J., & Muchenje, V. (2018). Multi-functional application of Moringa oleifera Lam. in nutrition and animal food products: A review. *Food research international*, 106, 317-334.
- Fidrianny, I., Kanapa, I., & Singgih, M. (2021). Phytochemistry and pharmacology of moringa tree: an overview. *Biointerface Res Appl Chem*, 11(3), 10776-10789.
- Fuglie, L. J. (1999). The miracle tree: Moringa oleifera, natural nutrition for the tropics.
- Kashyap, P., Kumar, S., Riar, C. S., Jindal, N., Baniwal, P., Guiné, R. P., Correia, P. M., Mehra, R., & Kumar, H. (2022). Recent advances in Drumstick (Moringa oleifera) leaves bioactive compounds: Composition, health benefits, bioaccessibility, and dietary applications. *Antioxidants*, 11(2), 402.
- Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., & Ogwal-Okeng, J. W. (2010). Phytochemicals and uses of Moringa oleifera leaves in Ugandan rural communities.
- Leone, A., Fiorillo, G., Criscuoli, F., Ravasenghi, S., Santagostini, L., Fico, G., Spadafranca, A., Battezzati, A., Schiraldi, A., & Pozzi, F. (2015). Nutritional characterization and phenolic profiling of Moringa oleifera leaves grown in Chad, Sahrawi Refugee Camps, and Haiti. *International journal of molecular sciences*, 16(8), 18923-18937.
- Luqman, S., Srivastava, S., Kumar, R., Maurya, A. K., & Chanda, D. (2012). Experimental assessment of Moringa oleifera leaf and fruit for its antistress, antioxidant, and scavenging potential using in vitro and in vivo assays. *Evidence-Based Complementary and Alternative Medicine*, 2012.
- Madziga, H., Sanni, S., & Sandabe, U. (2010). Phytochemical and elemental analysis of Acalypha wilkesiana leaf. *Journal of American Science*, 6(11), 510-514.
- Maheshwari, K., Yadav, R., Malhotra, J., Dhawan, N., & Mohan, L. (2014). Fascinating nutritional, prophylactic, therapeutic and socio-economic reconcile attributable to drum stick tree (Moringa oleifera Lam.). *Glob. J. Med. Res. B: Pharm. Drug Discov. Toxicol. Med*, 14, 11-22.
- Mbikay, M. (2012). Therapeutic potential of Moringa oleifera leaves in chronic hyperglycemia and dyslipidemia: a review. *Frontiers in pharmacology*, 3, 24.
- Mehra, R., Kumar, H., Kumar, N., & Kaushik, R. (2020). Red rice conjugated with barley and rhododendron extracts for new variant of beer. *Journal of Food Science and Technology*, 57, 4152-4159.
- Mir, M. A., Sawhney, S., & Jassal, M. (2013). Qualitative and quantitative analysis of phytochemicals of Taraxacum officinale. *Wudpecker Journal of Pharmacy and Pharmacology*, 2(1), 1-5.

- Padayachee, B., & Baijnath, H. (2020). An updated comprehensive review of the medicinal, phytochemical and pharmacological properties of *Moringa oleifera*. *South African Journal of Botany*, 129, 304-316.
- Patel, P., Patel, N., Patel, D., Desai, S., & Meshram, D. (2014). Phytochemical analysis and antifungal activity of *Moringa oleifera*. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(5), 144-147.
- Raj, A. J., Gopalakrishnan, V. K., Yadav, S. A., & Dorairaj, S. (2011). Antimicrobial activity of *Moringa oleifera* (Lam.) root extract. *Journal of Pharmacy Research*, 4(5), 1426-1427.
- Rios, J.-L., & Recio, M. C. (2005). Medicinal plants and antimicrobial activity. *Journal of ethnopharmacology*, 100(1-2), 80-84.
- Saadabi, A. M., & Zaid, I. A. (2011). An in vitro antimicrobial activity of *Moringa oleifera* L. seed extracts against different groups of microorganisms. *Australian Journal of Basic and Applied Sciences*, 5(5), 129-134.
- Sahakitpichan, P., Mahidol, C., Disadee, W., Ruchirawat, S., & Kanchanapoom, T. (2011). Unusual glycosides of pyrrole alkaloid and 4'-hydroxyphenylethanamide from leaves of *Moringa oleifera*. *Phytochemistry*, 72(8), 791-795.
- Sofowora, A. (1993). Medicinal plants and traditional medicine in Africa. Spectrum Books Limited. Ibadan, Nigeria, 1-153.
- Stevens, G., Baiyeri, K., & Akinnnagbe, O. (2013). Ethno-medicinal and culinary uses of *Moringa oleifera* Lam. in Nigeria. *Journal of medicinal plants research*, 7(13), 799-804.
- Sultana, B., Anwar, F., & Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*, 14(6), 2167-2180.
- Tshabalala, T., Ndhlala, A., Ncube, B., Abdelgadir, H., & Van Staden, J. (2020). Potential substitution of the root with the leaf in the use of *Moringa oleifera* for antimicrobial, antidiabetic and antioxidant properties. *South African Journal of Botany*, 129, 106-112.
- Verma, A. R., Vijayakumar, M., Mathela, C. S., & Rao, C. V. (2009). In vitro and in vivo antioxidant properties of different fractions of *Moringa oleifera* leaves. *Food and Chemical Toxicology*, 47(9), 2196-2201.
- Vongsak, B., Sithisarn, P., & Gritsanapan, W. (2012). HPLC quantitative analysis of three major antioxidative components of *Moringa oleifera* leaf extracts. *Planta Medica*, 78(11), PJ15.