

A preliminary recording of insects on the island of Farwa Northwest of Libya

Hoda M. Elmareme¹, Abdulhamed M. Etriiki^{2*}, Asmaa D. Mkhebes³, Ali A. Bataw⁴, Munay A. Albarbar⁵, Nahad S. Ben Omar⁶

Corresponding author: abdulhamed.etriiki@gu.edu.ly, Department of Zoology, Faculty of Science, University of Gharyan, Gharyan, Libya

¹ Department of Zoology, Faculty of Science, University of Tripoli, Tripoli, Libya.

³ Department of Ecology, Faculty of Science, Sabratha University, Sabratha, Libya

⁴ Department of Zoology, Faculty of Science, University of Omar Al Mukhtar, Al-Bayda, Libya

⁵ Department of Zoology, Faculty of Science, University of Tripoli, Tripoli, Libya.

⁶ Department of Zoology, Faculty of Science, University of Tripoli, Tripoli, Libya.

Received:
28 June 2024

Accepted:
30 August 2024

Publish online:
31 August 2024

Abstract

The study aimed to present the first record of insect families in Farwa Island. Farwa Island is the largest and most important island in Libya; located in Northern west part of Libya in the Mediterranean, its insect fauna is poorly studied. Short time intensive collection of insect samples was conducted in four areas (mouth of the valley, bay of containers, center of the Island, and Ras-Attalgha) in February, March and April 2021. Different methods were used for insect collection, sorting, preservation and later identification to the lowest taxonomic level and verification. 9 orders, 31 families, 40 genera and 58 insect species were collected in the study; the center of the Island had the highest numerical density and diversity of insect fauna due to its diverse vegetation compared to other areas. Diverse suitable habitats and absent insect control measures provided suitable environmental conditions for insects breeding, feeding and hiding from enemies. This study constitutes the first comprehensive survey of insect fauna on Farwa Island and adds significant value to the ongoing assessment of insect diversity in Libya. A long-term study has to be conducted to investigate detailed information about the abundance and diversity of insects and other arthropods on Farwa Island.

Keywords: Collection, Biodiversity, Farwa Island, Insect fauna, Libya

INTRODUCTION

Insects are invertebrate animals (Phylum: arthropods) that comprise the largest known and most widely distributed group (Class: Insecta) of multicellular organisms worldwide. Although the estimates of their numbers may vary, most entomologists agree that there are approximately 1.1 million described insect species. However, estimates of their actual number vary from 1.5 to 5 million species, and a consensus is converging on about 3.5 million species (roughly three times the number of described species) (Grimaldi, 2023). Insects are found in almost all environments, on plants, around



The Author(s) 2024. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, for non-commercial purposes only, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

buildings and under objects such as rocks and tree trunks; aquatic insects can be found in lakes, ponds, streams, rivers and swamps. Many insects have direct or indirect impacts on humans (pests as those considered parasitic, disease-transmitting vectors or homes/agricultural products destroying organisms). Moreover, insects are considered the most ecologically significant land organisms based on their large biomass. Some insect species are very destructive to forests and crops, but they constitute only a small percentage, and their numbers are biologically controlled by parasitoids and predatory insects. About 80-85% of the world's flowering plants are pollinated by insects; some insect products (honey, silk, dyes, wax and sustainable protein) are indispensable materials for humans. The fruit fly *Drosophila melanogaster* (as an experimental organism) has inspired the scientific society with many significant discoveries and breakthroughs in biology and other related fields (Grimaldi and Engel, 2005; Mohamed and Shaurub, 2010; Grimaldi, 2023).

Farwa Island is the biggest and most significant Island in Libya. It is situated 150 km west of Tripoli in the Northwest part of Libya on the Mediterranean Sea at the border region between Libya and Tunisia. The Island is roughly 12 km long with a maximum width of approximately 3 km, and the total surface area is about 31 km². Coordinates: 33°06'20.3"N 11°44'39.2"E. With its location in the Mediterranean climate zone, Farwa Island has a temperate climatic condition with sunny days and northern winds dominating most of the year. The mean annual temperature is 20°C and the mean annual rainfall reaches 190 mm. The Island is characterized by a diverse collection of habitats (beaches, drying salt lakes, trees, sand dunes, marshes, mudflats and widespread tidal areas). These diverse environments provide a suitable habitat for many floral and faunal species (Etayeb, 2002; Etayeb and Essghaier, 2007; EGA-RAC/SPA, 2012; Isemmann *et al.*, 2016).

Studies in Farwa Island on invertebrates groups generally and insect fauna particularly are poor. To the best of our knowledge, one previous study was conducted on invertebrates on this Island; the study by Jdeidi *et al.* (2015) revealed insects as the most abundant organisms (most of which were Coleoptera followed by Lepidoptera, Diptera, Hymenoptera and Orthoptera); it also included spiders and scorpions. This research introduces the first thorough study of recorded insect families in Farwa Island.

MATERIALS AND METHODS

The study was conducted during the months of February, March and April 2021. The Island was divided into four areas: mouth of the valley, bay of containers, center of the Island, and Ras-Attalgha. (Figure 1).

The following tools and equipment were used: tweezers, manual collection net, white cloths (2 x 3 meters), umbrella with a diameter of 1 meter, jars of different sizes, plastic/glass tubes and containers for collecting ground insects, magnifying glass, notebook for recording sample data (sample number, date of collection and place of sampling). Each area was visited where samples were collected using different collection methods from using forceps to methods of hitting and shaking branches of the bushes.

The samples were collected over two periods from 10 am to 1pm and from 2 pm to 5 pm, kept alive in jars or plastic/glass tubes, sorted and labeled in the field based on areas and deposited in the laboratory of the biology department, faculty of science, University of Tripoli. They were then killed using pieces of cotton filled with 75% ethanol in sealed jars/tubes, and then pinned in tightly sealed insect boxes for later identification and verification. Classification to the lowest possible taxonomic level was conducted at the University of Tripoli and laboratory of biology department, faculty of science, Omar Al-Mukhtar University (Al Bayda) using binocular microscope and special references (Zavattari, 1934; Borror *et al.*, 1981; Beccaloni *et al.*, 2005; Lamas, 2008; Massa, 2009;

Kirk-Spriggs and Sinclair, 2017). The data of each sample and the level of its classification were placed on special sample cards for the purpose of permanent preservation and reference when needed.




Figure: (1). Study area (Farwà Island) in the Northwest of Libya, divided into four regions: 1. mouth of the valley, 2. bay of containers, 3. center of the Island, and 4. Ras-Attalgha

RESULTS

The study collected 58 insect species belonging to 40 genera, 31 families and 9 orders, and they were arranged in (Table 1), followed by the scientific names for each insect. Four samples could be identified only to the family level (Asilidae, Reduvidae, Silphidae and Tetrigidae).

Order	Family	Scientific name
Lepidoptera Linnaeus 1758	Lycaenidae Leach 1815	<i>Aricia agestis</i> Denis and Schiffermüller 1775
	Noctuidae Latreille 1809	<i>Agrotis infusa</i> Boisduval 1832
	Pieridae Swainson, 1820	<i>Pieris brassicae</i> Linnaeus 1758
		<i>Pieris rapae</i> Linnaeus 1758
		<i>Pontia daplidice</i> Linnaeus 1758
Odonata Fabricius 1793	Pterophoridae Zeller 1841	<i>Gillmeria pallidactyla</i> Haworth 1811
	Zygaenidae Latreille 1809	<i>Zygaena felix</i> Oberthür 1876
	Libellulidae Rambur 1842	<i>Sympetrum decoloratum</i> Selys 1884
		<i>Sympetrum fonscolombii</i> Selys 1840
		<i>Acrotylus insubricus</i> Scopoli 1786
Orthoptera Latreille 1793	Acrididae MacLeay 1819	<i>Oedipoda caerulescens</i> Linnaeus 1758
		<i>Oedipoda fuscocincta</i> Lucas 1849
		<i>Oedipoda miniata</i> Pallas 1771
	Gryllotalpidae Saussure 1870	<i>Omocestus rufipes</i> Zetterstedt 1821
		<i>Truxalis nasuta</i> Linnaeus 1758
Thysanura Linnaeus 1758	Tetrigidae Rambur 1838	<i>Neocurtilla hexadactyla</i> Perty 1832
	Lepismatidae Linnaeus 1758	<i>Lepisma saccharina</i> Linnaeus 1758

DISCUSSION

Insects are indispensable organisms for life on earth and for the functioning and sustainability of all ecosystems (Saxena and Omkar, 2023). Based on a short-term intensive survey, this paper constitutes the first comprehensive collection of insects on Farwa Island. Moreover, in view of the importance of the study of insects in Libya in general and Farwa Island in particular, this study (represented by 9 orders, 31 families, 40 genera and 58 insect species) was compared to other similar studies; a study conducted by Yahiya (2014) included similar families (Coreidae, Lycaenidae, Lygaeidae, Pentatomidae, Pieridae). Similarly, a study by Mifsud (2000) had Carabidae, Coccinellidae, Formicidae, Sarcophagida and Tenebrionidae, and another study by EL-Meghrabi (2009) included Lygaeidae, Pentatomidae and Reduvidae, all of which were recorded in this survey. This study recorded the presence of families Zygaenidae, Noctuidae, Silphidae, Scoliidae, which were not present in the aforementioned studies (Yahiya, 2014), (Mifsud, 2000), (El-Meghrabi, 2009). 77 families obtained in the aforementioned studies were not recorded in this survey. This study was the first of its kind in the sampling and classification of insects on the Island, specifically among the areas and environments in which insects were collected and classified.

This insect biodiversity was due to the presence of many suitable habitats that may provide appropriate environments for both insects breeding and hiding from enemies. Moreover, the absence of control measures also helped to increase the abundance and distribution of insects in the study area

The results were obtained from samples collected only from Farwa Island and a multiplicity of insect habitats was observed, especially in the central region of the Island because of the diversity of vegetation compared to the number of visits to the study areas. On the other hand, the rest of the study areas, such as the mouth of the valley, and Ras-Attalgha, had the least numeri-

cal density and diversity of insect families due to the large number of waves and sea currents that result in strong winds for the two areas and thus the inappropriateness of climatic conditions, in addition to the scarcity of vegetation. The presence of insect families near plants was more abundant, followed by sand, and the least was near dung and carrion.

CONCLUSION

The results of this short time intensive survey of insects on Farwa Island showed that the Island has relatively abundant and diverse insect fauna. This is due to the fact that the Island provides suitable environments for insects breeding, feeding and escaping enemies; the absence of insect control measures adds to this rich biodiversity. Based on this comprehensive collection of insects from Farwa Island, this research adds important value to the continuously growing evaluation of the insect fauna in Libya.

ACKNOWLEDGEMENT

Authors declare there is no acknowledgement. No financial assistance was received in support of the study.

ETHICS

This article is original. The authors declare no ethical issues that may arise after the publication of this manuscript.

Duality of interest: The authors declare that there is no duality of interest associated with this manuscript.

Author contributions: All the authors contributed equally to preparing, developing and carrying out this manuscript.

Funding: This work did not receive any funding (institutional, private or corporate financial support). The authors self-funded their study.

REFERENCES

- Beccaloni G., Scoble M., Robinson G. & Pitkin, B. (2005) The Global Lepidoptera names index. Natural History Museum. Available from <http://www.nhm.ac.uk/our-science/data/lepindex/index.html>
- Borror, D.J., De Long, D.M., Triplehorn, C.A., (1981). An Introduction to the Study of Insects, fifth ed. Saunders College Publishing
- EGA-RAC/SPA Waterbird Census Team, (2012). Atlas of wintering water birds of Libya, 2005-2010. Imprimerie COTI, Tunisia.
- El-Meghrabi, M. S. (2009). Survey of some Heteropterous insects in Libya. Journal of Productivity and Development, 14(3), 747-758.
- Etayeb K.S., (2002). A study of Migratory and resident Marine Birds in Ras Attalgha and the Western part of Farwa island. M. Sc. Thesis. Tripoli University. Tripoli, Libya.

- Etayeb K.S. & Essghaier M.F.A., (2007). Breeding of marine birds on Farwa Island, western Libya. *Ostrich*, 78(2), 419–421. <http://dx.doi.org/10.2989/OSTRICH.78.2.48.128>.
- Grimaldi, D and Engel, M.S. (2005). *Evolution of the Insects*. Cambridge University Press, Cambridge, UK.
- Grimaldi, D. A. (Ed.). (2023). *The Complete Insect: Anatomy, Physiology, Evolution, and Ecology*. Princeton University Press.
- Isemmann P., Hering J., Brehme S., Essghaer M., Etayeb K.S., Bourass E. & Azafzaf H., (2016). *Birds of Libya*. Groupe Horizon, France, 302 pp.
- Jdeidi, T. B., Saed, F. A., and Elhosk, M. A. (2015). *The Main Fauna and Flora of Farwa Island, Libya*.
- Kirk-Spriggs, A.H. and Sinclair, B.J. (eds). (2017). *Manual of Afrotropical Diptera*. Volume 1. Introductory chapters and keys to Diptera families. Suricata 4. South African National Biodiversity Institute, Pretoria; pp. 267–355
- Lamas, G. (2008). Global butterfly names: various checklists. Taxome Project. Available from <http://www.ucl.ac.uk/taxome/gbn/>
- Linnaeus, C., (1758). *Systema Naturae, sive regna tria naturae systematice proposita per classes, ordines, genera et species*. Holmiae (Stockholm), 824pp.
- Massa, B. (2009). Annotated check-list of Orthoptera of Libya. *Journal of Orthoptera Research*, 18(1), 75–93. <http://dx.doi.org/10.1665/034.018.0109>
- Mifsud, D. (2000). Present knowledge of the entomofauna of the Maltese Islands. *Entomologica Basiliensia* 22: 75-86.
- Mohamed, W.F and Shaurub, E.S.H. (2010). A checklist of some recorded insects in Misurata, Libya. *Journal of King Saud University (Science)*, 22(2), 61-65. doi:10.1016/j.jksus.2010.02.001
- Saxena, S., Omkar (2023). *Insects in Scientific Research Advancement*. In: Omkar (eds) *Insects as Service Providers*. Springer, Singapore. https://doi.org/10.1007/978-981-19-3406-3_9
- Yahiya, Y.M, (2014). Common insects of orders Coleoptera, Lepidoptera and Hemiptera in middle region of Al-Jabal Al-Khdar, Libya. *Middle East Journal of Agriculture Research*, 3(3), 395-397.
- Zavattari, E. (1934). *Prodromo della fauna della Libia*. Tipografia già Cooperativa, Pavia, i-viii+1234pp