

# Bacteriological and statistical analysis of the causes behind urinary tract infections and the trends in antibiotic resistance in Ajdabiya.

*by* Husam Dakheel

---

**Submission date:** 05-Feb-2026 04:17AM (UTC-0600)

**Submission ID:** 2871668434

**File name:** ections\_and\_the\_trends\_in\_antibiotic\_resistance\_in\_Ajdabiya..doc (422.5K)

**Word count:** 3276

**Character count:** 19504

**Bacteriological and statistical analysis of the causes behind urinary tract infections and the trends in antibiotic resistance in Ajdabiya.**Husam Dakheel<sup>1\*</sup>

\*Corresponding author:  
[husam.alamamy@gmail.com](mailto:husam.alamamy@gmail.com)  
 Department of Biology, Faculty of Science, University of Ajdabiya, Ajdabiya, Libya

Received:

.....

Accepted:

.....

Publish online:

.....

**Abstract**

This study aimed to assess the prevalence of bacteria causing urinary tract infections (UTIs) and to identify common patterns of bacterial isolates along with their relationship to urine pH in patients visiting the Elite Clinic laboratory in Ajdabiya. A total of 60 urine samples were collected under sterile conditions from individuals suspected of having UTIs. Bacterial cultures were conducted using standard media, and the bacterial species were identified based on their growth characteristics and biochemical tests. Additionally, urine pH was measured during the analysis. The findings revealed that 48 samples (80%) showed positive bacterial growth, indicating a high incidence of UTIs. Gram-negative bacteria accounted for most isolates (70–75%), while Gram-positive bacteria represented a smaller proportion (25–30%). Among the identified pathogens, *Escherichia coli* was the most prevalent (40–45%), followed by *Klebsiella* spp. (15–20%) and other species at lower frequencies. Furthermore, most positive samples were associated with acidic urine conditions, with 60–65% of elevated bacterial growth observed at pH levels below 7, compared to 35–40% in alkaline urine. The study highlights a significant prevalence of UTIs in the area under investigation, with a predominance of Gram-negative bacteria, particularly *E.coli*. It underscores the influence of urine pH on bacterial growth and distribution, offering valuable insights into the dynamics of urinary pathogens.

**Keywords:** Urinary tract infections, urine culture, *Escherichia coli*, bacterial resistance, antibiotics.

**INTRODUCTION**

Urinary tract infections (UTIs) rank among the most prevalent bacterial illnesses encountered in clinical practice, affecting individuals across various age groups (Kundu et al., 2025; He et al., 2025; Baimakhanova et al., 2025). However, the incidence of these infections is notably higher in females than in males, primarily due to distinct anatomical and physiological differences. The widespread occurrence poses considerable public health concern, underscoring their significance within medical care (Li et al., 2022; Yang et al., 2022; Aleinein et al., 2024).

It may develop into serious complications if not diagnosed and treated properly, such as kidney inflammation or kidney failure (Scherberich et al., 2021; Kaur al., 2021; Mancuso al., 2023).



The Author(s) 2024. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

<sup>15</sup> Bacteria are the primary cause of urinary tract infections, with *E.coli* being the most common pathogen. Other types, including Klebsiella, Proteus, and Staphylococcus, are also significant contributors (Ahmed et al., 2014; Jalil et al., 2022; Zhou et al., 2023).

Understanding these causes has become increasingly critical in recent years, largely due to the steady rise in antibiotic resistance. This issue is widely recognized as one of the most urgent challenges in global health, especially given the widespread problem of indiscriminate or unregulated use of medications (Sabtu et al., 2015; Larsson and Flach, 2022; Salam et al., 2023).

This study is essential as it concentrates on local bacterial patterns responsible for urinary tract infections and evaluates their reactions to commonly prescribed antibiotics. Its goal is to support evidence-based treatment decisions, ultimately reducing treatment failures and curbing the spread of antibiotic resistance. Furthermore, the findings offer a credible scientific foundation for enhancing treatment and prevention strategies within healthcare institutions.

### <sup>13</sup> MATERIALS AND METHODS

This study was carried out at the Elite Clinic laboratory in Ajdabiya between September and December 2025. to investigate urinary tract infections among the city's residents. Employing a descriptive analytical approach, the research focused on bacteriological and statistical analysis of urine samples to determine the primary causes of infections, examine antibiotic resistance patterns, and identify the demographics most vulnerable to these infections. A total of 60 urine samples were collected from patients presenting to the clinic with clinical symptoms suggestive of a potential urinary tract infection. Participants included both genders and various age groups. Samples were obtained using the midstream collection method in sterile containers, following clear instructions given to patients on how to properly collect their samples to minimize contamination risks. They were transported to the laboratory within a 30-minute window after collection to maintain testing accuracy. Samples that arrived late or showed signs of contamination were excluded from the analysis.

#### 1.Laboratory tests

Laboratory tests were conducted following standard approved methods and included the following procedures:

- Microscopic analysis of urine to identify white blood cells, red blood cells, and bacteria.
- Culturing of urine samples on blood agar medium, with plates incubated at 37°C for 24 to 48 hours.
- Identification of isolated bacteria based on colony morphology, Gram staining, and specific biochemical tests tailored to each bacterial species .

#### 2.Antibiotic Sensitivity Testing

The susceptibility of bacterial isolates to frequently used antibiotics was evaluated through the Kirby-Bauer Disk Diffusion Method. Based on the diameters of the inhibition zones, the results were categorized as sensitive, intermediate, or resistant.

#### 3.Statistical analysis

The collected data were entered into organized tables to apply the analysis using descriptive statistical methods. The main objective was to determine percentages and frequencies, with the aim of clarifying the distribution of bacterium types and studying their patterns of resistance to antibiotics, in addition to analyzing the relationships between the variables studied. All statistical analyses were performed using Microsoft Excel 2021 and Python 3.11.

## RESULTS

### 1. General distribution of bacterial isolates

Urine culture results showed positive bacterial growth in 48 samples, representing 80% of the to-

tal samples studied, indicating a high prevalence of urinary tract infections among patients attending the Elite Clinic in Ajdabiya. Bacterial isolates varied between Gram-negative and Gram-positive bacteria, with a clear predominance of Gram-negative bacteria. The results showed that *E. coli* was the most common among the isolates, accounting for 31.3% (15 cases out of 48), followed by *Streptococcus* at 20.8% (10 cases), then *Candida* at 14.6% (7 cases). *Pseudomonas* and *Staphylococcus* were equally prevalent at 12.5% (6 cases each), while *Proteus mirabilis* was the least prevalent at 8.3% (4 cases) for example Figure 1. These results demonstrate the key role of *E. coli* as the main cause of urinary tract infections, due to its virulence factors that enable it to adhere to urinary epithelial cells and colonize the urinary tract.

The results also showed a difference in infection rates among different groups, with the highest infection rate recorded among males at 50%, followed by females at 43.3%, while the infection rate among children was the lowest, reaching 6.7% of the cases studied for example Figure 2. The higher incidence rate among females compared to adult males can be explained by anatomical and physiological factors such as the shortness of the urethra and its proximity to the anus, which facilitates the transmission of bacteria and increases the likelihood of infection.




Figure: (1). Distribution of Bacterial Isolates in Positive Urine Cultures




Figure: (2). Distribution of Urinary Tract Infection Cases by Gender

The prevalence of bacterial species found in urine samples, categorized by urine pH for example Figure 3. A noticeable variation in the occurrence of several bacterial species was seen between acidic and alkaline urine, suggesting that urine pH might influence bacterial proliferation and serious infection.




Figure: (3). Distribution of Urinary Tract Infection Cases by Gender

**2. Antibiotic sensitivity test results**

The sensitivity test results revealed significant variations in how bacterial isolates responded to the antibiotics tested, with differing resistance levels observed across the drugs. Augmentin and Amikacin showed the highest resistance rates at 17.9% each, suggesting a decline in their therapeutic effectiveness against certain bacterial strains. Moderate resistance was identified as Ceftriaxone, Azithromycin, Cefuroxime, and Amoxicillin, each with a resistance rate of 7.1%. In contrast, nalidixic acid exhibited a lower resistance rate of 5.4%, while gentamicin demonstrated the least resistance at 1.8%, highlighting its comparatively high efficacy against the isolates for example Figure 4 . These findings point to the prevalence of multidrug resistance, particularly among Gram-negative bacteria, which poses an increasing challenge for treatment. They emphasize the importance of relying on urine culture results and antibiotic susceptibility testing to guide the selection of appropriate therapies.




Figure: (4). Antibiotic resistance rates among isolated bacteria

## DISCUSSION

The present study demonstrated a high prevalence of urinary tract infections (UTIs), as evidenced by the high rate of positive urine cultures (80%). This finding indicates that UTIs represents a significant health concern among patients attending Al-Nokhba Clinic in Ajdabiya and is comparable to rates reported in similar regional and international studies (Khan et al., 2023; Srour et al., 2025).

Gram-negative bacteria predominate among the isolated pathogens, with *E.coli* being the most frequently identified organism. This result is consistent with numerous previous studies that identify *E. coli* as the primary etiological agent of UTIs worldwide (Alhazmi et al., 2023; Farag et al., 2024; El-Arifi et al., 2024; Wireko et al., 2025).

The predominance of *E. coli* is largely attributed to its virulence factors, including adhesins and fimbriae, which enhance attachment to uroepithelial cells and promote colonization of the urinary tract (Sung et al., 2024).

Other bacterial and fungal isolates such as *Streptococcus* spp, *Candida* spp, *Pseudomonas* spp, and *Staphylococcus* spp were detected at variable frequencies, suggesting possible roles of host immunity and the production of toxins, prior antibiotic exposure, and healthcare-associated factors (Bunduki et al., 2021; Wang et al., 2025; Gao et al., 2026).

Differences in infection rates were observed among gender groups, with males showing a slightly higher prevalence than females, while children accounted for the lowest proportion of cases. Although UTIs are typically more common in females due to anatomical and physiological factors, the higher rate among males in this study may be related to age distribution, urological abnormalities, or underlying conditions such as prostatic disorders (Brandstrom and Hansson, 2022; Gonzalez et al., 2024; Moghnia et al., 2025).

Variation in bacterial distribution according to urine pH was also observed, indicating that urinary acidity or alkalinity may influence bacterial growth and infection severity, as previously reported (Herrera-Espejo et al., 2024; Erinmez and Ozturk, 2024).

Antimicrobial susceptibility testing revealed variable resistance patterns, with the highest resistance

observed against Augmentin and Amikacin, while Gentamicin showed the lowest resistance rate. These findings indicate the presence of multidrug-resistant organisms, particularly among Gram-negative bacteria, emphasizing the necessity of routine urine culture and antibiotic susceptibility testing prior to treatment (Adjei, S and Adjei, 2025).

## CONCLUSION

This research determined that urinary tract infections constitute a notable health issue among attendees of the Elite Clinic in Ajdabiya, where urine culture outcomes showed a substantial rise in the proportion of affirmative samples, mirroring the broad presence of this affliction in the studied populace. The findings demonstrated that Gram-negative microbes were the primary driver of contagion, with *E. coli* being the most frequently found among bacterial separations, which align with observations from numerous prior investigations. The study additionally uncovered an interesting variety in bacterial isolates and pointed to variations in contagion frequency based on sex and age bracket, affirming the impact of anatomical and physiological aspects and existing medical ailments on infection initiation. Moreover, the evaluation emphasized that urine acidity can influence the proliferation and spread of certain microbial types, which is directly linked to the infection's intensity and the nature of the germ.

Concerning antimicrobial susceptibility assessment, the data exhibited a variability in how bacterial isolates reacted to the medications employed, with differing resistance profiles noted, particularly among Gram-negative bacteria, while certain antimicrobials displayed comparatively greater effectiveness. These outcomes suggest a worsening trend in microbial resistance to drugs, underscoring the need to utilize urine culture findings and sensitivity assays as the foundation for selecting suitable therapy.

The study suggests enhancing health awareness initiatives and encouraging the prudent use of antibiotics, alongside ensuring precise laboratory diagnoses. These strategies are intended to curb the spread of urinary tract infections and mitigate bacterial resistance, ultimately contributing to more effective treatments and safeguarding public health.

## ACKNOWLEDGEMENT

I would like to express his sincere thanks and gratitude to Hawa for her significant contribution and outstanding efforts in analyzing the samples, which played a key role in completing this scientific work accurately and efficiently.

He would also like to extend his sincere thanks to Samar for her valuable assistance and fruitful cooperation, which contributed to facilitating the work and completing it as required.

We would also like to express our deep gratitude and appreciation to the Elite Medical Center for providing us with the opportunity to work in its laboratories and for providing all the necessary facilities and equipment, which had a significant impact on the success of this research.

## ETHICS

The author declared that this study was conducted in compliance with ethical standards.

**Duality of interest:** The authors declare that they have/ have no duality of interest associated with this manuscript.

**Author contributions:** Husam Dakheel contributed to all aspects of the research and manuscript preparation.

**Funding:** This research was conducted without any external funding and was supported solely by the authors' own efforts.

## REFERENCES

- Adjei, S. K., & Adjei, P. (2025). Antimicrobial Susceptibility Patterns of Uropathogens: A Retrospective Study at a Primary Care Hospital in Ghana. *Health Science Reports*, 8(7), e71029.
- Ahmed, E., Hussain, M. S., Sheikh, S. I., & Ejaz, M. (2014). Urinary tract bacterial pathogens and their sensitivity pattern. *Journal of Rawalpindi Medical College*, 18(2).
- Aleinein, I. A., & Sokhn, E. S. (2024). Knowledge and prevalence of urinary tract infection among pregnant women in Lebanon. *Heliyon*, 10(17).
- Alhazmi, A. H., Alameer, K. M., Abuageelah, B. M., Alharbi, R. H., Mobarki, M., Musawi, S., & Dhayhi, N. (2023). Epidemiology and antimicrobial resistance patterns of urinary tract infections: a cross-sectional study from Southwestern Saudi Arabia. *Medicina*, 59(8), 1411.
- Baimakhanova, B., Sadanov, A., Trenozhnikova, L., Balgimbaeva, A., Baimakhanova, G., Orasymbet, S., & Turgumbayeva, A. (2025). Understanding the Burden and Management of Urinary Tract Infections in Women. *Diseases*, 13(2), 59.
- Brandstrom, P., & Hansson, S. (2022). Urinary tract infection in children. *Pediatric Clinics*, 69(6), 1099-1114.
- Bunduki, G. K., Heinz, E., Phiri, V. S., Noah, P., Feasey, N., & Musaya, J. (2021). Virulence factors and antimicrobial resistance of uropathogenic *Escherichia coli* (UPEC) isolated from urinary tract infections: a systematic review and meta-analysis. *BMC infectious diseases*, 21(1), 753.
- El-Arifi, S., AbdAlla, E., Mahgoub, E. S., Fadol, B., & Elmubarak, R. (2024). Characteristics and antibiotic resistance patterns of urinary tract isolates in hospitalized and non-hospitalized patients: a cross-sectional study in Khartoum, Sudan. *BMC Infectious Diseases*, 24(1), 1356.
- Erinmez, M., & Ozturk, M. (2026). Urinary Tract Infection in Patients with Urolithiasis: A Large Retrospective Observational Study of Clinical Features and Microbiological Spectrum. *Pathogens*, 15(1), 98.
- Farag, P. F., Albulushi, H. O., Eskembaji, M. H., Habash, M. F., Malki, M. S., Albadrani, M. S., & Hanafy, A. M. (2024). Prevalence and antibiotic resistance profile of UTI-causing uropathogenic bacteria in diabetics and non-diabetics at the Maternity and Children Hospital in Jeddah, Saudi Arabia. *Frontiers in Microbiology*, 15, 1507505.
- Gao, J., Wu, F., Zhang, Y., Wu, J., Li, W., Su, N., & Wang, Y. N. (2026). Drug Resistance Patterns and Pathogen Distribution in Patients with Urinary Tract Infection in Suzhou, China: A Five-Year Retrospective Study (2020–2024). *Infection and Drug Resistance*, 1-15.

- Gonzalez, M. J., Robino, L., Zunino, P., & Scavone, P. (2024). Urinary tract infection: is it time for a new approach considering a gender perspective and new microbial advances?. *Frontiers in Urology*, 4, 1487858.
- He, Y., Zhao, J., Wang, L., Han, C., Yan, R., Zhu, P., & He, W. (2025). Epidemiological trends and predictions of urinary tract infections in the global burden of disease study 2021. *Scientific Reports*, 15(1), 4702.
- Herrera-Espejo, S., Domínguez-Miranda, J. L., Rodríguez-Mogollo, J. I., Pachón, J., Cordero, E., & Pachón-Ibáñez, M. E. (2024). Effects of pH on the Pathogenicity of *Escherichia coli* and *Klebsiella pneumoniae* on the Kidney: In Vitro and In Vivo Studies. *International Journal of Molecular Sciences*, 25(14), 7925.
- Jalil, M. B., & Al Atbee, M. Y. N. (2022). The prevalence of multiple drug resistance *Escherichia coli* and *Klebsiella pneumoniae* isolated from patients with urinary tract infections. *Journal of clinical laboratory analysis*, 36(9), e24619.
- Kaur, R., & Kaur, R. (2021). Symptoms, risk factors, diagnosis and treatment of urinary tract infections. *Postgraduate medical journal*, 97(1154), 803-812.
- Khan, M., Hussain, F., Naseem, M., Clementina, R., Gul, N., Habib, A., & Khan, I. (2024). Investigating Urinary Pathogens and Their Antibiotic Resistance: A Cross-Sectional Urine Culture Study. *Cureus*, 16(7).
- Kundu, N., Mou, F. A., Samia, F. N., Urmi, C. Y., Hossain, M. S., & Ahmad, T. (2025). Knowledge-level, attitudinal-trend and preventive-practices on urinary tract infections among reproductive-aged female university students. *Discover Public Health*, 22(1), 587.
- Larsson, D. J., & Flach, C. F. (2022). Antibiotic resistance in the environment. *Nature Reviews Microbiology*, 20(5), 257-269.
- Li, X., Fan, H., Zi, H., Hu, H., Li, B., Huang, J., & Zeng, X. (2022). Global and regional burden of bacterial antimicrobial resistance in urinary tract infections in 2019. *Journal of clinical medicine*, 11(10), 2817.
- Mancuso, G., Midiri, A., Gerace, E., Marra, M., Zummo, S., & Biondo, C. (2023). Urinary tract infections: the current scenario and future prospects. *Pathogens*, 12(4), 623.
- Moghnia, O. H., Al Otaibi, H. S., Al Haqqan, A. M., Sokhn, E. S., Pathan, S. S., Abdulaziz, N. E., & Al-Sweih, N. A. (2025). Prevalence of Urinary Tract Infections and Antibiotic Susceptibility Patterns of Uropathogens among Neonates in Maternity Hospital, Kuwait: A Six-Year Retrospective Study. *Medical Principles and Practice*, 34(3), 250-261.
- Sabtu, N., Enoch, D. A., & Brown, N. M. (2015). Antibiotic resistance: what, why, where, when and how?. *British medical bulletin*, 116(1), 105-113.
- Salam, M. A., Al-Amin, M. Y., Salam, M. T., Pawar, J. S., Akhter, N., Rabaan, A. A., & Alqumber, M. A. (2023, July). Antimicrobial resistance: a growing serious threat for global public health. In *Healthcare* (Vol. 11, No. 13, p. 1946). MDPI.

- Scherberich, J. E., Funfstuck, R., & Naber, K. G. (2021). Urinary tract infections in patients with renal insufficiency and dialysis—epidemiology, pathogenesis, clinical symptoms, diagnosis and treatment. *GMS infectious diseases*, 9, Doc07.
- Sror, N., Bornstein, G., Golan, L., Rimbrot, D., Barlev, Y., Freund, O., & Zornitzki, L. (2025). Rate and Predictors of Urine Culture Positivity Following Single-Dose Antibiotic Administration: a Prospective Single-Center Study. *Infectious Diseases and Therapy*, 1-13.
- Sung, K., Nawaz, M., Park, M., Chon, J., Khan, S. A., Alotaibi, K., & Khan, A. A. (2024). Comprehensive genomic analysis of uropathogenic *E. coli*: virulence factors, antimicrobial resistance, and mobile genetic elements. *Pathogens*, 13(9), 794. Challenges and Prospects. *Microorganisms*, 13(12), 2714.
- Wang, B., Wang, Y., Liu, H., Yu, M., Wang, S., Liu, L., & Tan, H. (2025). Current Scientific Advances in Vaccines Against UTIs:
- Wireko, S., Ngmaakan, G., Dwomoh, D., Hansen, C., Nartey, S., & Senu, E. (2025). Regional Burden of Urinary Tract Infection, Its Aetiologies, and Antibiotic Resistance Patterns, 2020–2023: A Retrospective Study. *Health Science Reports*, 8(11), e71503.
- Yang, X., Chen, H., Zheng, Y., Qu, S., Wang, H., & Yi, F. (2022). Disease burden and long-term trends of urinary tract infections: A worldwide report. *Frontiers in public health*, 10, 888205.
- Zhou, Y., Zhou, Z., Zheng, L., Gong, Z., Li, Y., Jin, Y., & Chi, M. (2023). Urinary tract infections caused by uropathogenic *Escherichia coli*: mechanisms of infection and treatment options. *International journal of molecular sciences*, 24(13), 10537.

**Author’s Contribution Form**

Please Tick  in the appropriate box for the author:

| S.N | Author’s Name*       | Con-ceived& design of the study     | Collect-ed the data                 | Contributed data or anal-ysis tools | Performed the analy-sis/ inter-pretation of data | Drafting the manu-script/revising forimportant in-tellectual con-text | Approval of the final version of the manu-script** |
|-----|----------------------|-------------------------------------|-------------------------------------|-------------------------------------|--|---|--|
| 1   | <b>Husam Dakheel</b> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/>              | <input checked="" type="checkbox"/>                                   | <input checked="" type="checkbox"/>                |
| 2   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |
| 3   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |
| 4   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |
| 5   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |
| 6   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |
| 7   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |
| 8   | —                    | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>            | <input type="checkbox"/>                         | <input type="checkbox"/>  | <input type="checkbox"/>                           |

Other contribution:

| Author’s number | Specify contribution in more detail ( no more than one sentence)   |
|-----------------|--|
| 1               | The author was responsible for all aspects of the study, including study execution, data validation, and final submission. |
| 2               | —  |
| 3               | —  |
| 4               | —  |
| 5               | —  |
| 6               | —  |
| 7               | —  |

| S.N | Author’s name | Email                   | Author’s signature      |
|-----|---------------|-------------------------|-------------------------|
| 1   | Husam Dakheel | husam.alamamy@gmail.com | <i>Husam S. Dakheel</i> |

|   |   |   |   |
|---|---|---|---|
| 2 | — | — | — |
| 3 | — | — | — |
| 4 | — | — | — |
| 5 | — | — | — |
| 6 | — | — | — |
| 7 | — | — | — |

# Bacteriological and statistical analysis of the causes behind urinary tract infections and the trends in antibiotic resistance in Ajdabiya.

## ORIGINALITY REPORT

|                  |                  |              |                |
|------------------|------------------|--------------|----------------|
| 10%              | 6%               | 7%           | 0%             |
| SIMILARITY INDEX | INTERNET SOURCES | PUBLICATIONS | STUDENT PAPERS |

## PRIMARY SOURCES

|   |  |    |
|---|--|----|
| 1 | <a href="https://journals.su.edu.ye">journals.su.edu.ye</a><br>Internet Source   | 1% |
| 2 | Manjuri Sharma, Faheem Nazir Qanoongo, Prodip Kumar Doley, Gayatri Pegu, Miranda Pegu. "Spectrum Of Urinary Tract Infection Among Renal Allograft Recipients in a Tertiary Care Centre of Northeast India", Springer Science and Business Media LLC, 2024<br>Publication   | 1% |
| 3 | Iman Abu Aleinein, Elie Salem Sokhn. "Knowledge and prevalence of urinary tract infection among pregnant women in Lebanon", Heliyon, 2024<br>Publication   | 1% |
| 4 | Al-Mufarjiyah, Aisha Said Salim. "Uncomplicated Urinary Tract Infection in Oman: A Prospective Insight of the Etiology, Resistome and Diagnostic Antimicrobial Stewardship", Sultan Qaboos University (Oman), 2025<br>Publication  | 1% |
| 5 | Hooraf Afshar Naderi, Farimah Beheshti, Ahad Ali Harati Asl, Vida Hojati, Omid Azizi. "Chronic uropathogenic Escherichia coli infections in adolescence: in vitro analysis of clinical isolates, multidrug resistance, and in vivo hippocampal cognitive impacts", Naunyn- | 1% |

# Schmiedeberg's Archives of Pharmacology, 2025

Publication

---

|    |  |      |
|----|--|------|
| 6  | <a href="https://docs.neu.edu.tr">docs.neu.edu.tr</a><br>Internet Source   | 1 %  |
| 7  | <a href="https://journals.abu.edu.ng">journals.abu.edu.ng</a><br>Internet Source   | 1 %  |
| 8  | <a href="http://www.ijrar.org">www.ijrar.org</a><br>Internet Source  | <1 % |
| 9  | <a href="http://www.ijsr.net">www.ijsr.net</a><br>Internet Source  | <1 % |
| 10 | <a href="http://phr.emrystick.com">phr.emrystick.com</a><br>Internet Source  | <1 % |
| 11 | DiMaggio, Dennis Anthony, Jr.. "Investigating the Regulation of Secreted Virulence Factors by the Sae Two-Component System in Staphylococcus aureus", Georgetown University, 2025<br>Publication | <1 % |
| 12 | Wen-Ta Chiu, Jonathan Wu. "Decoding the Pandemic - Unveiling the Hidden Depths of the COVID-19 Iceberg", Routledge, 2025<br>Publication  | <1 % |
| 13 | <a href="http://agribalkan.congress.gen.tr">agribalkan.congress.gen.tr</a><br>Internet Source  | <1 % |
| 14 | <a href="http://bau.edu.lb">bau.edu.lb</a><br>Internet Source  | <1 % |
| 15 | <a href="http://epu.edu.iq">epu.edu.iq</a><br>Internet Source  | <1 % |
| 16 | Avalos, Alexis A.. "Deciphering the Source of Escherichia coli in Urinary Tract Infections: A Comparative Metagenomic Analysis of the  | <1 % |

# Female Urogenital Microbiome.", Loyola University Chicago

Publication

---

**17** Tahir, Bilal. "The Efficacy of Pakistan's National Action Plan in Addressing Antimicrobial Resistance", Hamad Bin Khalifa University (Qatar), 2024 <1 %

Publication

---

**18** [etd.aau.edu.et](http://etd.aau.edu.et) <1 %

Internet Source

---

**19** [orbi.uliege.be](http://orbi.uliege.be) <1 %

Internet Source

---

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On